



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## GEOMETRY.

**510. Proposed by JOSEPH E. ROWE, State College, Pa.**

Show how to find the equation of a line parallel to a side of the triangle of reference and passing through a given point, in any system of homogeneous coördinates, using the condition that two lines are parallel in this system but not the condition that two lines are perpendicular. Illustrate the method by using it to find the trilinear coördinates of the points of contact of one escribed circle of the triangle.

**511. Proposed by FRANK V. MORLEY, Student, Haverford College, Pa.**

Let  $a_i$  ( $i = 1, 2, 3, 4$ ) be four points on a circle and let the in-center of the triangle formed by omitting  $a_i$  be  $c_i$ ; prove that the four points  $c_i$  form a rectangle.

## CALCULUS.

**425. Proposed by O. S. ADAMS, U. S. Coast and Geodetic Survey, Washington, D. C.**

Show that the infinite product

$$(1 - z)(1 + \tfrac{1}{2}z)(1 - \tfrac{1}{3}z)(1 + \tfrac{1}{4}z) \cdots = \frac{\sqrt{\pi}}{\Gamma(1 + \tfrac{1}{2}z)(\tfrac{1}{2} - \tfrac{1}{2}z)}.$$

**426. Proposed by C. N. SCHMALL, New York City.**

If  $A$  be the area of a plane triangle constructed with the sides  $a, b, c$ , such that

$$a^3 + b^3 + c^3 = 3k^3,$$

show that the maximum value of  $A$  is  $\frac{1}{4}k^2$ .

## MECHANICS.

**342. Proposed by WILLIAM HOOVER, Columbus, Ohio.**

A uniform rod of length  $2a$  is freely hinged at one end, at the other end a string of length  $b$  is attached which is fastened at its further end to a point on the surface of a homogeneous sphere of radius  $c$ . If the masses of the rod and sphere are equal, find the motion of the system when slightly disturbed from the vertical, and the cubic equation giving the corresponding small oscillations.

**343. Proposed by J. ROSENBAUM, New Haven, Conn.**

Two bodies of equal masses and coefficients of friction  $\mu_1$  and  $\mu_2$  are connected by a light spring of stiffness  $k$  and placed on an inclined plane. Discuss the motion of each body when the angle between the non-stretched spring and the plane is  $\theta$ .

## NUMBER THEORY.

**261. Proposed by NORMAN ANNING, Chilliwack, B. C.**

Show that for any positive integer  $n$  (excluding powers of 2) positive integers  $a_1, a_2, a_3, \dots, a_k$ , which are less than  $n/2$  can be chosen in such a way that

$$2^k \cos(a_1\pi/n) \cos(a_2\pi/n) \cos(a_3\pi/n) \cdots \cos(a_k\pi/n) = 1.$$

**262. Proposed by C. N. SCHMALL, New York City.**

If  $x, y, z$  are three integers, consecutive among the integers prime to 3, show that

$$x(x - 2y) - z(z - 2y) = \pm 3.$$

## SOLUTIONS OF PROBLEMS.

## ALGEBRA.

J. A. BULLARD and J. W. BALDWIN solved 464. These solutions were received after selections for publication were made.

**465. Proposed by CYRUS B. HALDEMAN, Ross, Ohio.**

Having given  $\tan^{-1} 1 = \tan^{-1} 1/2 + \tan^{-1} 1/3$ , show that

$$\tan^{-1} 1 = 5 \tan^{-1} 1/8 + 2 \tan^{-1} 1/18 + 3 \tan^{-1} 1/57.$$

SOLUTION BY HORACE OLSON, Chicago, Illinois.

Put  $\tan^{-1} 1/2 = \tan^{-1} 1/3 + \tan^{-1} x_1 = \tan^{-1} \frac{3x_1 + 1}{3 - x_1}$ . From this equation  $x_1$  is found to be  $1/7$ . Hence,  $\tan^{-1} 1 = 2 \tan^{-1} 1/3 + \tan^{-1} 1/7$ .

Now put

$$\tan^{-1} 1/3 = \tan^{-1} 1/7 + \tan^{-1} x_2 = \tan^{-1} \frac{7x_2 + 1}{7 - x_2};$$

whence  $x_2 = 2/11$ , and  $\tan^{-1} 1 = 3 \tan^{-1} 1/7 + 2 \tan^{-1} 2/11$ .

Now put

$$\tan^{-1} 1/7 = \tan^{-1} x_3 + \tan^{-1} y_3 = \tan^{-1} \frac{x_3 + y_3}{1 - x_3 y_3}.$$

This gives the indeterminate equation  $x_3 y_3 + 7(x_3 + y_3) = 1$ , or  $(x_3 + 7)(y_3 + 7) = 50$ . One set of solutions of this equation is  $x_3 = 1/8$ ,  $y_3 = 1/57$ . Hence,  $\tan^{-1} 1 = 3 \tan^{-1} 1/8 + 3 \tan^{-1} 1/57 + 2 \tan^{-1} 2/11$ . Put

$$\tan^{-1} 2/11 = \tan^{-1} 1/8 + \tan^{-1} x_4 = \tan^{-1} \frac{8x_4 + 1}{8 - x_4};$$

whence  $x_4 = 1/18$ . Hence,  $\tan^{-1} 1 = 5 \tan^{-1} 1/8 + 2 \tan^{-1} 1/18 + 3 \tan^{-1} 1/57$ .

Also solved by E. B. ESCOTT, W. J. THOME, M. T. REED, G. W. HARTWELL, and E. E. WHITEFORD.

**466. Proposed by E. B. ESCOTT, Kansas City, Mo.**

For what functions,  $f$ , are the following relations true:

When

$$\frac{f(x, y, z)}{X} = \frac{f(y, z, x)}{Y} = \frac{f(z, x, y)}{Z},$$

then

$$\frac{f(X, Y, Z)}{x} = \frac{f(Y, Z, X)}{y} = \frac{f(Z, X, Y)}{z}?$$

SOLUTION BY ALBERT A. BENNETT, University of Texas.

We notice that  $f(X, Y, Z)$  must be homogeneous, since if we replace  $X, Y, Z$  by  $cX, cY, cZ$ , in the first relation it is unaltered, and hence also the second. Thus  $x, y, z$  may be regarded as the homogeneous coordinates of the points in one plane,  $X, Y, Z$  of those in a second. The problem may therefore be expressed as follows: What are the involutonic plane transformations with triangular symmetry? Here "involutonic" is used in the restricted sense as of period two, and triangular symmetry is used in approximately the sense first suggested by Clifford, the point  $(1, 1, 1)$  being a center of triangular symmetry. There can be little doubt that there exist transformations of this form which are essentially transcendental and which may be regarded as limiting cases of algebraic birational transformations. To catalogue the explicit forms of even the algebraic cases is perhaps out of the question, since no explicit classification of algebraic forms of Cremona transformations has been attempted beyond the very simplest cases. Normal forms under the Cremona group are indeed known. Compare Pascal's *Repertorium* or other detailed articles on birational geometry.

It is furthermore obvious that a given geometric solution gives rise to an infinite number of analytic solutions. For example, if  $g(x, y, z)$  be one solution, and  $\varphi(x, y, z)$  be a constant or the equation of any triangularly symmetric self-corresponding curve whether the self-correspondence be singular or not, then  $\varphi(x, y, z)g(x, y, z)$  is a solution. Apart from the trivial solutions  $f(x, y, z) = 0$  and  $f(x, y, z) = x$ , the simplest case is probably the quadratic transformation given by  $f(x, y, z) = 1/x$ , and then taking  $\varphi(x, y, z) = xyz$ , or  $\sqrt{xyz}$ , or  $xyz/\log(1 + xyz)$  or  $(xy + yz + zx)(x + y + z)$ , etc., we get other related solutions from  $f(x, y, z) = 1/x$ .